

CLAIMS:

1. A method of making a no wash bead based assay, the method comprising:

preparing a first reagent comprising a buffer;

preparing a second reagent comprising a protein;

preparing beads of preselected size and having a coefficient of variation less than 5%, including washing the beads in the buffer to form a bead-buffer matrix and reducing the surfactancy of the beads to an effective amount;

adding an antigen for detecting the presence of a target species to the bead-buffer matrix such that the antigen attaches to the beads to form a bead-antigen mixture, the surfactancy of the beads facilitating attachment of the antigen thereto;

adding buffer to the bead-antigen mixture and thereafter incubating the mixture; and

adding second reagent to the bead-antigen mixture to reduce or eliminate non-specific binding sites.

2. A method as claimed in claim 1 wherein the first reagent is a carbonate buffer.

3. A method as claimed in claim 2 wherein the carbonate buffer

has a pH in the range of 9.0 - 10.0.

4. A method as claimed in claim 3 wherein the carbonate buffer has a pH of 9.6.

5

5. A method as claimed in claim 1 wherein the second reagent is bovine serum albumin (BSA).

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000

6. A method as claimed in claim 5 wherein the BSA comprises a 0.1 - 5.0% BSA in saline.

7. A method as claimed in claim 6 wherein the BSA is a 0.5% BSA in saline.

8. A method as claimed in claim 1 wherein the size of the beads is selected from one or more of the group consisting of 3 $\mu$  latex beads, 4 $\mu$  latex beads, 5 $\mu$  latex beads, 6 $\mu$  latex beads, 7 $\mu$  latex beads, 8 $\mu$  latex beads, 9 $\mu$  latex beads and 10 $\mu$  latex beads.

9. A method as claimed in claim 8 wherein the beads are selected so as to have a coefficient of variation not exceeding 5%.

10. A method as claimed in claim 9 wherein the beads are selected so as to have a coefficient of variation not exceeding 1.3%.

11. A method as claimed in claim 8 wherein multiple sizes of beads are selected.

12. A method as claimed in claim 1 wherein the antigens added are selected from the group consisting of RNP/SM, SM, SS-A, SS-B, SCL-70 and dsDNA.

13. A method as claimed in claim 1 wherein the antigens are selected from one of more of the group consisting of histones, lipids, viral antibodies, viral antigens, bacterial antibodies, bacterial antigens, recombinant proteins, and cellular antigens.

14. A method as claimed in claim 1 wherein the surfactancy of the beads is reduced to no more than 5% in order to enhance the ability to coat the beads with antigens.

15. A method as claimed in claim 14 wherein the surfactancy is no more than 0.5% of the beads.

16. A method as claimed in claim 1 wherein the bead-based assay is prepared in a flat-bottom container.

17. A method as claimed in claim 1 wherein the bead-buffer matrix is subjected to at least one prewashing step.

18. A method as claimed in claim 1 further comprising the step of centrifuging the bead-buffer matrix and the bead-antigen mixture, and the resuspension thereof.

19. A method as claimed in claim 1 further comprising the step of vortexing the bead-buffer matrix and the bead-antigen mixture, and the resuspension thereof.

20. A method of manufacturing a no wash kit for carrying out a bead based assay for testing for the presence of a target substance, the method comprising:

preparing a first reagent comprising a buffer;

preparing a second reagent comprising a protein;

preparing a third reagent comprising an indicator antibody, the third reagent being selected for its ability to identify the target substance;

preparing beads of preselected size and having a coefficient of variation less than 5%, including washing the beads in the buffer to form a bead-buffer matrix and reducing the surfactancy of the beads to an effective amount;

5        adding an antigen for detecting the presence of the target substance to the bead-buffer matrix such that the antigen attaches to the beads to form a bead-antigen mixture, the surfactancy of the beads facilitating attachment of the antigen thereto;

10        adding buffer to the bead-antigen mixture and thereafter incubating the mixture;

105873866-060401        adding second reagent to the bead-antigen mixture to reduce or eliminate non-specific binding sites;

      placing the bead-antigen mixture in a first container for use in an assay test procedure; and

15        placing the third reagent in a second container for use during the test procedure, the third reagent being used after the target substance has attached to the bead for the purpose of identifying the target substance.

20        21. A method as claimed in claim 20 wherein the third reagent comprises goat anti-human Ig.

22. A method as claimed in claim 20 comprising a plurality of bead-antigen mixtures, wherein a plurality of different antigens are each attached to a bead of different size.

23. A method as claimed in claim 22 wherein the bead sizes are selected from the following: 3 $\mu$  latex beads, 4 $\mu$  latex beads, 5 $\mu$  latex beads, 6 $\mu$  latex beads, 7 $\mu$  latex beads, 8 $\mu$  latex beads, 9 $\mu$  latex beads and 10 $\mu$  latex beads.

24. A method as claimed in claim 22 wherein one or more of the antigens are selected from the group consisting of RNP/SM, SM, SS-A, SS-B, SCL-70 and dsDNA.

25. A no wash kit for carrying out a bead based assay for testing for the presence of a target substance, the kit comprising a bead based assay prepared according to the method of claim 1, and a detector assay reagent comprising an indicator antibody, the detector assay reagent being selected for its ability to identify the target substance.

26. A no wash bead based assay for testing for the presence of a target substance, the assay being prepared according to the method

of claim 1.

09873866-060401  
FOUO "986E/850